

The effects of Martian near surface conditions on the photochemistry of amino acids

Inge Loes ten Kate^{a,b,*}, James R.C. Garry^a, Zan Peeters^a, Bernard Foing^c,
Pascale Ehrenfreund^a

^a*Leiden Institute of Chemistry, Astrobiology Laboratory, P.O. Box 9502, 2300 RA Leiden, The Netherlands*

^b*Leiden Observatory, Leiden University, P.O. Box 9513, 2300 RA Leiden, The Netherlands*

^c*ESA Research and Scientific Support Department, ESTEC/SCI-SR, P.O. Box 299, 2200 AG Noordwijk, The Netherlands*

Received 24 August 2005; accepted 2 December 2005

Available online 19 January 2006

Abstract

In order to understand the complex multi-parameter system of destruction of organic material on the surface of Mars, step-by-step laboratory simulations of processes occurring on the surface of Mars are necessary. This paper describes the measured effects of two parameters, a CO₂ atmosphere and low temperature, on the destruction rate of amino acids when irradiated with Mars-like ultraviolet light (UV). The results show that the presence of a 7 mbar CO₂ atmosphere does not affect the destruction rate of glycine, and that cooling the sample to 210 K (average Mars temperature) lowers the destruction rate by a factor of 7. The decrease in the destruction rate of glycine by cooling the sample is thought to be predominantly caused by the slower reaction kinetics. When these results are scaled to Martian lighting conditions, cold thin films of glycine are assumed to have half-lives of 250 h under noontime peak illumination. It has been hypothesised that the absence of detectable native organic material in the Martian regolith points to the presence of oxidising agents. Some of these agents might form via the interaction of UV with compounds in the atmosphere. Water, although a trace component of Mars' atmosphere, is suggested to be a significant source of oxidising species. However, gaseous CO₂ or adsorbed H₂O layers do not influence the photodestruction of amino acids significantly in the absence of reactive soil. Other mechanisms such as chemical processes in the Martian regolith need to be effective for rapid organic destruction.

© 2005 Elsevier Ltd. All rights reserved.

Keywords: Mars; Amino acids; Water; Photochemistry

1. Introduction

Mars is the target for future space missions with the search for traces of extinct or extant life as one of the main goals. Recent space missions have provided a wealth of information about the surface and atmospheric conditions on Mars (e.g. Squyres et al., 2004a, b; Formisano et al., 2004). The Martian atmosphere is dominated by CO₂ (95%). Other major gases in Mars' atmosphere are (percentage by moles) nitrogen (N₂, 2.7%), argon (Ar, 1.6%), oxygen (O₂, 0.13%) and carbon monoxide (CO,

0.08%). Water (H₂O) is a minor constituent (varying between 10 and 1000 parts per million (ppm), Encrenaz et al., 2004a), as well as methane (CH₄, 5 parts per billion (ppb), Formisano et al., 2004; Krasnopolsky et al., 2004). The fractional abundance of O₃ in the Martian atmosphere, with a mixing ratio of 10⁻⁸, is directly related to the level of O₂, while the abundance of O₂ is regulated by catalytic cycles involving HO_x species. Hydrogen peroxide (H₂O₂) has been suggested as a possible oxidising agent of the Martian surface. Photochemical models suggest that the mean column density of H₂O₂ should be in the range 10¹⁵–10¹⁶ cm⁻², and that H₂O₂ and H₂O abundances should be correlated. Encrenaz et al. (2004b) report an H₂O₂ atmospheric mixing ratio of 5 × 10⁻⁸ around the sub-solar point.

*Corresponding author. Astrobiology Laboratory, Leiden Institute of Chemistry, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands. Tel.: +31 71 5274541; fax: +31 71 5274397.

E-mail address: tenkate@strw.leidenuniv.nl (I.L. ten Kate).

No organic matter was detected in Martian regolith samples analysed by the two Viking landers (Biemann et al., 1977). Given the current presumed rate of meteoritic infall, detectable amounts of organic matter should have accumulated at the surface of Mars (Flynn and McKay, 1988). To account for this discrepancy, the effects of UV (ten Kate et al., 2005), gaseous oxidants (Oró and Holzer, 1979) and surface catalysts (Quinn and Zent, 1999) on organic molecules have been examined in laboratory experiments. Of particular interest is the stability of simple amino acids, as they may be important biomarkers searched for by instruments on the payload of future missions to Mars.

When irradiated with Mars-like UV, 300 nm thick polycrystalline films of amino acids degrade with half-lives of around 2×10^6 s (glycine) and 3×10^5 s (D-alanine) (ten Kate et al., 2005). However, this destruction was measured exclusively at room temperature and in vacuum ($\sim 10^{-6}$ mbar). In the current paper, we have examined the photostability of glycine samples at low temperature or in the presence of CO₂. These conditions are representative of the equatorial regions of Mars (~ 210 K, Kieffer et al., 1992) with a low adsorbed water content (Möhlmann, 2002) and no condensed CO₂. Regolith mineralogy and chemistry are not taken into account in these experiments. The experiments and the simulation chamber used for these experiments are described in Section 2, where a description of the used UV source is also given. In Section 3, we report on the infra-red (IR) spectroscopy of the irradiated glycine samples, and we tabulate the half-lives and UV destruction cross-sections measured in the reported experiments. The implications of the obtained results are explained in detail in Section 4.

2. Equipment and experimental procedure

2.1. Equipment and experiments

Glycine (99.7% purity, Merck) was deposited on silicon substrates, to form films with a thickness of 300 ± 50 nm, using a vacuum sublimation system, details of which are given in ten Kate et al. (2005). The films produced in this system have been shown to be optically thin (ten Kate et al., 2005). Experiments have been performed in a modified version of the system as used by Peeters et al. (2003) (Fig. 1). The system consists of a small vacuum chamber equipped with several gas-inlet ports. A closed-cycle, two-stage helium cryostat (Air products, Displex DE-202), which can be rotated while maintaining the vacuum, allows the temperature of the sample to be controlled to within 0.3 K in the range 20–300 K. In this system a Mars-like atmosphere, ~ 7 mbar CO₂ (Praxair, 99.996%), was created at room temperature, representing a Mars-like daytime surface temperature in equatorial regions. Alternatively, samples were cooled down to 210 K under vacuum to simulate an average Mars surface temperature. The silicon substrate is mounted on the cryostat, allowing the vapour




Fig. 1. Cryosystem used for the irradiation of amino acids.

deposited amino acid sample layer to face either a UV source (see Section 2.2) or the beam of a Fourier transform IR spectrometer (Excalibur FTS-4000, BioRad, $4000\text{--}500\text{ cm}^{-1}$ at 4 cm^{-1} resolution). Simultaneous cooling and using a 7 mbar atmosphere of CO₂ was not possible to achieve in our cryosystem, because CO₂ would selectively freeze on parts of the cryostat that are far colder (~ 40 K) than the 210 K at which the substrate window was maintained.

2.2. UV source

The UV source is a deuterium discharge lamp (Heraeus-Noblelight, DX 202), which has been calibrated against a known UV standard (Bentham R48) using a monochromator equipped with a photomultiplier tube. The measured spectrum of the DX202 lamp is shown in Fig. 2, along with a curve showing a representative noontime UV spectrum for Mars that was calculated for a low atmospheric dust-load by Patel et al. (2002). The integrated flux in the wavelength range 190–325 nm (on the amino acid layer) is 1.2×10^{14} photons/cm² s in the current system. This is ~ 12 times lower than the integrated flux on Mars in the same wavelength range (1.4×10^{15} photons/cm² s).

As has been pointed out by Schuerger et al. (2003), a deuterium discharge light source is not an exact fit to the




Fig. 2. Lighting spectrum (a) experienced by the equatorial noontime surface of Mars at $L_S = 70^\circ$ and (b) of the deuterium lamp. The grey area marks the 190–220 nm range, which is used in the discussion.

lighting spectrum found at Mars' surface. The deuterium emission spectrum peaks at around 200 nm, and is deficient in longer wavelengths compared to the UV spectrum at the Martian surface. Thus, the work presented here places upper limits on the half-life of a sample under Martian conditions.

3. Results

Thin films of glycine deposited on silicon substrates have been irradiated with UV in vacuum ($\sim 10^{-7}$ mbar), in a CO_2 atmosphere (~ 7 mbar) and at a temperature of 210 K. Multiple experiments have been performed for each modified parameter to achieve high accuracy of the measurements. Fig. 3 shows the natural logarithm of the normalised integrated absorbance ($\ln(S/S_0)$) plotted against irradiation time for all experiments. The slope of the linear fit through $\ln(S/S_0)$ represents the destruction rate. Table 1 lists the temperature of the glycine samples, the average pressure in the system and the corresponding average destruction rate of the different experiments. Also listed are the half-lives when extrapolated to a Martian noontime equatorial UV flux of 1.4×10^{15} photons/s/cm². The destruction rates found in the vacuum-only experiments were in agreement with the values found by ten Kate et al. (2005), with a small shift within the error bars, due to the use of a different system.

The system can be heated under vacuum prior to the experiments so that water is desorbed from the stainless steel. The equilibrium amount of water accreted on the disc during 24 h cooling at 210 K was measured with the IR spectrometer to be of the order of 10^{16} mol/cm². The amount of water and other contaminants in the CO_2 gas is in the order of 1 ppm. The background pressure in the system was in the order of 10^{-7} mbar.




Fig. 3. Natural logarithm of the normalised integrated absorbance ($\ln(S/S_0)$) plotted against irradiation time, for the deuterium irradiation of glycine (A) in vacuum at 294 K, (B) in a 7 mbar CO_2 atmosphere at 294 K, and (C) in vacuum when cooled to 210 K. The different lines represent the destruction rates of the separate experiments that have been performed.

4. Discussion

The effects of low temperature or a CO_2 atmosphere on the photodestruction rate of amino acids have been examined. This has been done by irradiating 300 ± 50 nm thick polycrystalline films of glycine deposited on silicon discs in the presence of a CO_2 atmosphere or when cooled to 210 K, corresponding to an average temperature on the surface of Mars. The results show no measurable effect of a CO_2 atmosphere on the destruction of glycine. However, when the samples are cooled to 210 K the destruction rate decreases by a factor of 7.

4.1. Effect of a CO_2 atmosphere

Thin films of glycine deposited onto silicon substrates have been irradiated with UV in a CO_2 atmosphere

Table 1

Average destruction rates of thin films of glycine measured under different conditions in the laboratory when subjected to UV irradiation

Temperature (K)	Number of repeated experiments	Atmosphere (mbar)	Destruction rate J ($\times 10^{-7} \text{ s}^{-1}$)	Mars half-life (s)
294	3	$\sim 10^{-7}$	4.5 ± 2.3	$1.3 \pm 0.8 \times 10^5$
294	4	~ 7	4.9 ± 2.3	$1.3 \pm 0.6 \times 10^5$
210	2	$\sim 10^{-7}$	0.8 ± 2.9	$0.9 \pm 7.5 \times 10^6$

Also listed are the extrapolated half-lives of glycine for a noontime equatorial flux on Mars.

Table 2

Noontime solar UV flux at the surface of Mars, UV flux of the deuterium lamp and average absorption (σ_{abs}) and scattering (σ_{scat}) cross-sections of CO₂ for two wavelength ranges

Wavelength range (nm)	Mars flux ^c ($\times 10^{12}$ photon/cm ² s)	Lamp flux ^d ($\times 10^{13}$ photon/cm ² s)	σ_{abs} (cm ²)	σ_{scat} (cm ²)
190–203 ^a	4	1	3×10^{-23}	1×10^{-24}
203–220 ^b	10	2	7×10^{-25}	7×10^{-25}

^aShemansky (1972).

^bKaraïskou et al. (2004).

^cPatel et al. (2002).

^dMeasured from 200 nm and up, 190–200 nm adapted from the spectrum provided by Heraeus.

(~ 7 mbar). Adding a CO₂ atmosphere is assumed to have two effects—extinction of the UV flux reaching the glycine film, and formation of O-radicals by photodissociation of CO₂. The extinction of UV can be divided into an absorption and a Rayleigh scattering component, both with a corresponding cross-section (σ , cm²), σ_{abs} and σ_{scat} , respectively. The wavelength range displayed in Fig. 2 can be divided into three regions: <203, 203–220 and >220 nm. In the region >220 nm the UV extinction is dominated by scattering with a very small cross-section $\sigma_{\text{scat}} < 10^{-25}$ cm² (upper limit of 10^7 scattered or absorbed photons/s), and is therefore not further considered in this discussion. In the region 203–220 nm both scattering and absorption contribute equally to the UV extinction, while in the region 190–203 nm absorption is the dominant process (Karaïskou et al., 2004; Shemansky, 1972). Table 2 gives an overview of the solar noontime UV flux on the surface of Mars and the output of UV flux from the deuterium lamp, as well as the average absorption and scattering cross-sections of CO₂, for the 190–203 and 203–220 nm intervals.

Above 167 nm, the dissociation of CO₂ occurs only through the reaction



Between 190 and 200 nm the CO-formation quantum yield is ~ 1 , but above 200 nm this value decreases to 0.16 at 214 nm, and no photodissociation occurs above 227 nm (Okabe, 1978). The formation rate of O-radicals will therefore be dominated by UV in the wavelength range 190–200 nm. As can be seen from Table 2, the UV absorption of CO₂ between 190 and 200 nm is small,

leading to an upper limit on the O radical formation in the order of 10^9 s^{-1} .

In summary, the total UV extinction in our experiments, through scattering and absorption by CO₂ over the full range of our lamp (190–325 nm), is in the order of 10^9 photons/s. The formation rate of O-radicals by the photodissociation of CO₂ is $\sim 10^9 \text{ s}^{-1}$. These values are both a factor 10^5 lower than the UV flux produced by the deuterium lamp; thus, the destruction rate of glycine thin films is *expected* to be dominated by photodestruction by UV. This concurs with the experimental results, presented in Table 1, where we found no effect on the destruction rate of glycine, when a CO₂ atmosphere was added and noontime equatorial illumination on Mars was simulated.

4.2. Effect of cooling

Thin films of glycine deposited on silicon substrates have been irradiated with UV when cooled to 210 K. The cooling resulted in a lower destruction rate of glycine, which is expected from reaction kinetics. At 210 K water is accreted onto the glycine film, as has been shown by background water accretion measurements (see Section 3). Water accretion may influence the destruction rate; the reaction rate could either decrease due to UV absorption by the ice layer, or increase due to formation of, e.g. OH radicals.

The amount of water accreted on the sample during 24 h cooling, inferred from measurements made on a blank disc under the same conditions, is in the order of 10^{16} molecules/cm², corresponding to a layer thickness of ~ 5 H₂O molecules (~ 1.5 nm). Table 3 shows absorption coefficients of water (Thompson et al., 1963) and the corresponding

Table 3
Absorption coefficients of water and absorbance of a 1.5-nm-thick water layer

Wavelength (nm)	Absorption coefficient (cm ⁻¹)	Absorbance
190	2×10^{-1}	3×10^{-8}
200	3×10^{-3}	5×10^{-10}
250	1×10^{-4}	2×10^{-11}

absorbance of the water layer on our samples at several wavelengths. These absorbances imply that in the wavelength range emitted by the deuterium lamp (190–325 nm) the absorption of UV photons by H₂O molecules hardly plays a role. The decrease in the destruction rate of the glycine film by cooling the sample, as measured in our experiments, is therefore thought to be predominantly caused by the slower reaction kinetics. A model of the upper Martian surface from Möhlmann (2004) suggests an average of two monolayers at 200 K, adsorbed on the porous surface of Martian soil. If only photodissociation of water adsorbed on amino acid layers is taken into account, the effect of these H₂O monolayers on the surface of Mars is negligible.

4.3. Water vapour

It has been postulated that the action of energetic UV photons on the water vapour present in the Martian troposphere leads to the formation of OH radicals, which influence the photochemistry of the soil (Hunten, 1979). Above 200 nm, UV absorption by gaseous water is negligible, between 190 and 200 nm the absorption cross-section is smaller than 10^{-21} cm² (Chung et al., 2001; Parkinson and Yoshino, 2003). The amount of residual water present as contamination in the CO₂ atmosphere in the chamber is in the order of 10^{-3} mbar, comparable to the average amount of gas phase H₂O in the Martian atmosphere. The water vapour content in the atmosphere of Mars varies between 10 and 1000 ppm (Encrenaz et al., 2004a). The amount of water in our system leads to a total absorption of 10^9 UV photons in the duration of the experiment. This is less than the total UV absorption and scattering by the CO₂ gas present in the system (see Section 4.1). A control experiment has been performed in a different system (described in ten Kate et al., 2005), in which a thin film of glycine was irradiated in a 10 mbar atmosphere consisting of 50% CO₂ and 50% H₂O (data not shown here). The destruction rate found in this experiment was the same as the destruction rates of glycine thin films irradiated in vacuum. Photodestruction of H₂O into OH radicals does not occur at wavelengths above 190 nm (Okabe, 1978), so is not expected to play a role in our experiments. On Mars, H₂O will be dissociated efficiently only in upper atmospheric layers, where UV irradiation is not substantially attenuated. The diffusion of radicals, formed by dissociation of water, to the surface,

and their subsequent reactions are not well understood. UV irradiation of H₂O vapour close to the surface will not result in a significant amount of reactive species.

5. Conclusions

We measured the destruction rate of ~300-nm-thick polycrystalline films of glycine deposited on silicon substrates, when irradiated with UV (190–325 nm) in vacuum ($\sim 10^{-7}$ mbar), in a CO₂ atmosphere (~ 7 mbar), and when cooled to 210 K. Regolith mineralogy and chemistry are not taken into account in these experiments. The results show that the presence of a 7 mbar CO₂ atmosphere does not affect the destruction rate of glycine by UV; the extinction of UV (with an extinction rate of 10^9 photons/s) and the formation of O radicals (with a formation rate of 10^9 s⁻¹) are very small compared to the flux of the deuterium lamp. However, cooling the amino acid during irradiation reduces the destruction rate by a factor of 7. A thin layer of water (representative for Martian conditions, see Möhlmann, 2002, 2004) accreted on the glycine film did not measurably influence the destruction rate. When the results on thin films of glycine by ten Kate et al. (2005) and the results of this work are scaled for Martian noontime lighting conditions, glycine exposed to UV at temperatures between 210–295 K has a half-life of approximately 35–250 h under continuous irradiation. Irradiation of 24 h in our experiments corresponds to an irradiation time of ~2 h on the surface of Mars at noontime. Only in the polar regions, during their respective summers, may sunlight shine on the surface for periods of >24 h, however, with lower intensity than at the equator. The simple geometric relationship between latitude and surface intensity for a sphere allows one to scale the life time of glycine calculated for noon illumination conditions at the equator to other latitudes. Our low temperature experiments performed at 210 K are more representative of mid- and high-latitude regions on Mars, and indicate that in those environments the destruction rate of amino acids may be reduced further. Furthermore, the amount of water adsorbed on the surface has a regional, seasonal and diurnal dependence. Therefore, the actual life time of exposed glycine on the cold surface of Mars will depend on several parameters. Using the model of amino acids embedded in Martian regolith described by ten Kate et al. (2005), with a mixing ratio of glycine of 1 ppb, we find a half-life of glycine of $\sim 10^8$ yr.

Energetic UV photons are suggested to form OH radicals from the water vapour present in the Martian atmosphere (e.g. Nair et al., 1994; Atreya and Gu, 1995). This dissociation takes place most efficiently in upper atmospheric layers, where UV irradiation is less attenuated. The OH radicals formed in the atmosphere and other photochemical processes could play a role in the formation of oxidising agents in the regolith (Zent and McKay, 1994). However, OH radical production on the Martian surface can also occur through processes involving adsorbed H₂O,

Fe(II) ions and H₂O₂ (Southworth and Voelker, 2003). It has been suggested that water abundance plays an important role in controlling reaction kinetics by triggering oxidative reactions involving photochemically produced dry acids that are adsorbed onto the soil (Quinn et al., 2005). Interaction of UV with Mars-analogue minerals in the presence of an oxygen atmosphere with a partial pressure comparable to that in the Martian atmosphere showed the formation of oxygen radicals (O₂⁻) on the minerals that likely destroy organic material (Yen et al., 2000). The stability of the intrinsic organic component of Martian soil analogues has recently been investigated by Garry et al. (2006). Those results support the idea that the key element in the destruction of organic material is the interaction of accreted water with the soil in the presence of radiation.

In summary, we studied the behaviour of amino acids irradiated by 190 to 325 nm UV in a CO₂ atmosphere and when cooled to 210 K. Our results form a basis for the understanding of more complex processes occurring on the Martian surface, in the presence of regolith and other reactive agents. Future research should involve a diurnal UV irradiation and temperature cycle, enabling to simulate the diurnal water frost deposition on the surface. Low temperatures may enhance the stability of amino acids in certain cold habitable environments, which may be important in the context of the origin of life.

Acknowledgements

ILtK is supported by the BioScience Initiative of Leiden University; JG is supported by SRON National Institute for Space Research, grant MG-058; PE and ZP are supported by Grant NWO-VI 016.023.003. We acknowledge the support from ESA-ESTEC and SRON.

References

- Atreya, S.K., Gu, Z.G., 1995. The photochemistry and stability of the atmosphere of Mars. *Adv. Space Res.* 16 (6), 57–68.
- Biemann, K., Oró, J., Toulmin III., P., Orgel, L.E., Nier, A.O., Anderson, D.M., Simmonds, P.G., Flory, D., Diaz, A.V., Rushneck, D.R., Biller, J.E., Laffleur, A.L., 1977. The search for organic substances and inorganic volatile compounds in the surface of Mars. *J. Geophys. Res.* 82, 4641–4658.
- Chung, C.-Y., Chew, E.P., Cheng, B.-M., Bahoub, M., Leeb, Y.-P., 2001. Temperature dependence of absorption cross-section of H₂O, HOD, and D₂O in the spectral region 140–193 nm. *Nucl. Instrum. Meth. Phys. Res. A* 467–468, 1572–1576.
- Encrenaz, Th., Lellouch, E., Atreya, S.K., Wong, A.S., 2004a. Detectability of minor constituents in the Martian atmosphere by infrared and submillimeter spectroscopy. *Planet. Space Sci.* 52, 1023–1037.
- Encrenaz, Th., Bézard, B., Greathouse, T.K., Richter, M.J., Lacy, J.H., Atreya, S.K., Wong, A.S., Lebonnois, S., Lefèvre, F., Forget, F., 2004b. Hydrogen peroxide on Mars: evidence for spatial and seasonal variations. *Icarus* 170, 424–429.
- Flynn, G.J., McKay, D.S., 1988. Meteorites on Mars. In: Lunar and Planetary Institute, Workshop on Mars Sample Return Science, pp. 77–78.
- Formisano, V., Atreya, S., Encrenaz, Th., Ignatiev, N., Giuranna, M., 2004. Detection of methane in the atmosphere of Mars. *Science* 306, 1758–1761.
- Garry, J.R.C., ten Kate, I.L., Martins, Z., Nørnberg, P., Ehrenfreund, P., 2006. Analysis and survival of amino acids in Martian regolith analogues. *Meteoritics Planet. Sci.* (accepted for publication 28 October 2005).
- Hunten, D.M., 1979. Possible oxidant sources in the atmosphere and surface of Mars. *J. Mol. Evol.* 14, 71–78.
- Karaïskou, A., Vallance, C., Papadakis, V., Vardavas, I.M., Rakitzis, T.P., 2004. Absolute absorption cross-section measurements of CO₂ in the ultraviolet from 200 to 206 nm at 295 and 373 K. *Chem. Phys. Lett.* 400, 30–34.
- Kieffer, H.H., Jakosky, B.M., Snyder, C.M., 1992. The planet Mars: from antiquity to present. In: Kieffer, H.H., Jakosky, B.M., Snyder, C.W., Matthews, M.S. (Eds.), Mars. The University of Arizona Press.
- Krasnopolsky, V.A., Maillard, J.P., Owen, T.C., 2004. Detection of methane in the Martian atmosphere: evidence for life? *Icarus* 172, 537–547.
- Möhlmann, D., 2002. Adsorption water in mid- and low-latitude Martian soil. In: Sawaya-Lacoste, H. (Ed.), ESA SP-518 Proceedings of Second European Workshop on Exo-/Astro-Biology, Graz, first ed. European Space Agency, Noordwijk, pp. 169–172.
- Möhlmann, D., 2004. Water in the upper Martian surface at mid- and low-latitudes: presence, state, and consequences. *Icarus* 168, 318–323.
- Nair, H.M., Allen, M., Anbar, A.D., Yung, Y.L., Clancy, R.T., 1994. A photochemical model of the Martian atmosphere. *Icarus* 111, 124–150.
- Okabe, H., 1978. Photochemistry of triatomic molecules. In: Okabe, H. (Ed.), Photochemistry of Small Molecules. Wiley, New York.
- Oró, J., Holzer, G., 1979. The photolytic degradation and oxidation of organic compounds under simulated Martian conditions. *J. Mol. Evol.* 14 (1–3), 153–160.
- Parkinson, W.H., Yoshino, K., 2003. Absorption cross-section measurements of water vapor in the wavelength region 181–199 nm. *Chem. Phys.* 294, 31–35.
- Patel, M.R., Zarnecki, J.C., Catling, D.C., 2002. Ultraviolet radiation on the surface of Mars and the Beagle 2 UV sensor. *Planet. Space Sci.* 50 (9), 915–927.
- Peeters, Z., Botta, O., Charnley, S.B., Ruitkamp, R., Ehrenfreund, P., 2003. The astrobiology of nucleobases. *Astrophys. J.* 593 (2), L129–L132.
- Quinn, R.C., Zent, A.P., 1999. Peroxide-modified titanium dioxide: a chemical analog of putative Martian soil oxidants. *Origins Life Evol. Biosphere* 29 (1), 59–72.
- Quinn, R.C., Zent, A.P., Grunthaner, P., Taylor, C.L., Garry, J.R.C., 2005. Detection and characterization of oxidizing acids in the Atacama Desert using the Mars Oxidation Instrument. *Planet. Space Sci.* 53 (13), 1376–1388.
- Schuerger, A.C., Mancinelli, R.L., Kern, R.G., Rothschild, L.J., McKay, C.P., 2003. Survival of endospores of *Bacillus subtilis* on spacecraft surfaces under simulated Martian environments: implications for the forward contamination of Mars. *Icarus* 165, 253–276.
- Shemansky, D.E., 1972. CO₂ extinction coefficient 1700–3000 Å. *J. Chem. Phys.* 56 (4), 1582–1587.
- Southworth, B.A., Voelker, B.M., 2003. Hydroxyl radical production via the photo-Fenton reaction in the presence of fluvic acid. *Environ. Sci. Technol.* 37, 1130–1136.
- Squyres, S.W., Arvidson, R.E., Bell III., J.F., Brückner, J., Cabrol, N.A., Calvin, W., Carr, M.H., Christensen, P.R., Clark, B.C., Crumpler, L., Des Marais, D.J., d’Uston, C., Economou, T., Farmer, J., Farrand, W., Folkner, W., Golombek, M., Gorevan, S., Grant, J.A., Greeley, R., Grotzinger, J., Haskin, L., Herkenhoff, K.E., Hviid, S., Johnson, J., Klingelhöfer, G., Knoll, A.H., Landis, G., Lemmon, M., Li, R., Madsen, M.B., Malin, M.C., McLennan, S.M., McSween, H.Y., Ming, D.W., Moersch, J., Morris, R.V., Parker, T., Rice Jr., J.W., Richter, L., Rieder, R., Sims, M., Smith, M., Smith, P., Soderblom, L.A., Sullivan, R., Wänke, H., Wdowiak, T., Wolff, M., Yen, A., 2004a. The Spirit rover’s Athena science investigation at Gusev Crater. *Mars. Science* 305, 794–799.

- Squyres, S.W., Arvidson, R.E., Bell III, J.F., Brückner, J., Cabrol, N.A., Calvin, W., Carr, M.H., Christensen, P.R., Clark, B.C., Crumpler, L., Des Marais, D.J., d'Uston, C., Economou, T., Farmer, J., Farrand, W., Folkner, W., Golombek, M., Gorevan, S., Grant, J.A., Greeley, R., Grotzinger, J., Haskin, L., Herkenhoff, K.E., Hviid, S., Johnson, J., Klingelhöfer, G., Knoll, A.H., Landis, G., Lemmon, M., Li, R., Madsen, M.B., Malin, M.C., McLennan, S.M., McSween, H.Y., Ming, D.W., Moersch, J., Morris, R.V., Parker, T., Rice Jr., J.W., Richter, L., Rieder, R., Sims, M., Smith, M., Smith, P., Soderblom, L.A., Sullivan, R., Wänke, H., Wdowiak, T., Wolff, M., Yen, A., 2004b. The Opportunity rover's Athena science investigation at Meridiani Planum, Mars. *Science* 306, 1698–1703.
- ten Kate, I.L., Garry, J.R.C., Peeters, Z., Quinn, R., Foing, B., Ehrenfreund, P., 2005. Amino acid photostability on the Martian surface. *Meteoritics Planet. Sci.* 40 (5), 1185–1193.
- Thompson, B.A., Harteck, P., Reeves Jr., R.R., 1963. Ultraviolet absorption coefficients of CO₂, CO, O₂, H₂O, N₂O, NH₃, NO, SO₂, and CH₄ between 1850 and 4000 Å. *J. Geophys. Res.* 68 (24), 6431–6436.
- Yen, A.S., Kim, S.S., Hecht, M.H., Frant, M.S., Murray, B., 2000. Evidence that the reactivity of the Martian soil is due to superoxide ions. *Science* 289, 1909–1912.
- Zent, A.P., McKay, C.P., 1994. The chemical reactivity of the Martian soil and implication for future mission. *Icarus* 108, 146–157.